

A GEOMETRIC DESCRIPTION OF THE $\mathrm{PSL}_4(\mathbb{R})$ -HITCHIN COMPONENT

SAMUEL A. BALLAS

ABSTRACT. These notes form a rough outline of the correspondence between the $\mathrm{PSL}_4(\mathbb{R})$ -Hitchin component and convex foliated projective structures from [3].

1. INTRODUCTION

Let Σ be an orientable hyperbolic surface. The *Teichmüller space* of Σ , denoted $\mathcal{T}^2(\Sigma)$, consists of conjugacy classes of discrete and faithful representations from $\Gamma := \pi_1(\Sigma)$ into $\mathrm{PSL}_2(\mathbb{R})$. This set is well known to have the topology of a ball of dimension $3|\chi(\Sigma)|$. There is a well known unique irreducible representation, ρ_n , from $\mathrm{PSL}_2(\mathbb{R})$ into $\mathrm{PSL}_n(\mathbb{R})$ coming from the natural action of $\mathrm{PSL}_2(\mathbb{R})$ on the symmetric product, $\mathrm{Sym}^{n-1}(\mathbb{R}^2) \cong \mathbb{R}^n$. In his article [6], Hitchin showed that the the component of the space of conjugacy classes, $\mathfrak{X}(\Gamma, \mathrm{PSL}_n(\mathbb{R}))$, containing the image of $\mathcal{T}^2(\Sigma)$ under ρ_n is also a ball of dimension $(n^2 - 1)|\chi(\Sigma)|$. This component is typically referred to as the *Hitchin component* and we will denote it as $\mathcal{T}^n(\Sigma)$.

For small values of n these Hitchin component can be thought of as moduli spaces of geometric structures on manifolds. For $n = 2$, the Hitchin component is the same as Teichmüller space which allows us to identify $\mathcal{T}^2(\Sigma)$ with the space of marked hyperbolic structures on Σ . More specifically, let $[\rho] \in \mathcal{T}^2(\Sigma)$. Since $\mathrm{PSL}_2(\mathbb{R})$ can be identified with the orientation preserving isometry group of \mathbb{H}^2 we can identify the quotient $\mathbb{H}^2/\rho(\Gamma) \cong \Sigma$, thus giving a marked hyperbolic structure on Σ . Conversely, the holonomy representation of any hyperbolic structure is a discrete and faithful representation from Γ into $\mathrm{PSL}_2(\mathbb{R})$, and equivalent marked hyperbolic structures have conjugate holonomy.

When $n = 3$, work of Goldman [2] and Goldman-Choi [1] show that the space $\mathcal{T}^3(\Sigma)$ can be identified with the moduli space of marked convex projective structures on Σ . Roughly speaking, a convex projective structure on Σ is a realization of Σ as $\Omega/\rho(\Gamma)$, where Ω is a convex set that sits inside of an affine patch (see Exercise 1) of \mathbb{RP}^2 and $\rho : \Gamma \rightarrow \mathrm{PSL}_3(\mathbb{R})$ is a discrete and faithful representation. For example, when ρ factors through $\mathrm{PSL}_2(\mathbb{R})$ the set Ω can be taken to be the unit disk in $\mathbb{R}^2 \subset \mathbb{RP}^2$ (See Exercise 2). While it is easy to associate a conjugacy class of discrete faithful representations with a marked projective structure, it is much more difficult to start with a representation and find an appropriate convex projective structure.

Exercise 1. Let \mathbb{RP}^n be the quotient of $\mathbb{R}^{n+1} \setminus \{0\}$ by the action of \mathbb{R}^\times by scaling and let H be a hyperplane \mathbb{RP}^n . $\mathbb{RP}^n \setminus H$ is called an affine patch. The purpose of this exercise is to justify this terminology.

- (1) Show that an affine patch can be identified with the affine space \mathbb{R}^n .

- (2) Show that the subgroup $\mathrm{PGL}_{n+1}(\mathbb{R})$ consisting of elements that preserve H is equivariantly isomorphic (with respect to the identification from part (1)) to the affine group of matrices of the form

$$\begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix},$$

where $A \in \mathrm{GL}_n(\mathbb{R})$ and $b \in \mathbb{R}^n$.

Exercise 2. Let $\langle x, y \rangle = x_1y_1 + x_2y_2 - x_3y_3$ be the standard bilinear form of signature $(2, 1)$ on \mathbb{R}^3 . Let $C = \{x \in \mathbb{R}^3 \mid \langle x, x \rangle < 0\}$ be the cone of vectors with negative self pairing. The image, $P(C)$, of C in \mathbb{RP}^2 serves as a model for \mathbb{H}^2 .

- (1) Let H be the image of the x_1x_2 -plane in \mathbb{RP}^2 . Show that in the affine patch defined by H that $P(C)$ can be identified with the unit disk. This model is known as the Klein Model
- (2) Let K be the image of the plane $x_3 - x_2 = 0$ in \mathbb{RP}^2 . Show that in the affine patch defined by that $P(C)$ can be identified with the set $v > u^2$. (Hint, use coordinates $u = x_1, v = x_3 + x_2, w = x_3 - x_2$ and use inhomogeneous coordinates $w = 1$). This model is known as the paraboloid model.

Our goal in this lecture to explain a correspondence between the space $\mathcal{T}^4(\Sigma)$ and certain types of projective structures on the unit tangent bundle, $S\Sigma$, of Σ . This result is originally due to Guichard and Wienhard [3]. The rough idea is that there is an \mathbb{R} action on $S\Sigma$ given by the geodesic flow. This flow gives rise to a pair of foliations, \mathcal{F} and \mathcal{G} , which are referred to as the *stable foliation* and *geodesic foliation*, respectively. We will show that representations in $\mathcal{T}^4(\Sigma)$ correspond to projective structures in which these foliations can be realized in a geometrically meaningful way inside of \mathbb{RP}^3 . When $[\rho] \in \mathcal{T}^4(\Sigma)$ factors through $\mathrm{PSL}_2(\mathbb{R})$ then we can regard these projective structures as projective realizations of the familiar $\widetilde{\mathrm{SL}}_2(\mathbb{R})$ structures on these manifolds.

2. $S\Sigma$ AND THE GEODESIC FLOW

In this section we will discuss important properties of $M := S\Sigma$ and the action of the geodesic flow. Let $\bar{\Gamma} = \pi_1(M)$, then $\bar{\Gamma}$ is a central (non-split) extension of Γ that fits into the following short exact sequence.

$$(2.1) \quad 0 \rightarrow \mathbb{Z} \rightarrow \bar{\Gamma} \rightarrow \Gamma \rightarrow 1.$$

The manifold M has an important regular cover, \bar{M} , corresponding from the \mathbb{Z} subgroup in (2.1). Furthermore, the foliations \mathcal{F} and \mathcal{G} lift to foliations $\bar{\mathcal{F}}$ and $\bar{\mathcal{G}}$ of \bar{M} . The foliations \mathcal{F} and \mathcal{G} can also be lifted to the universal cover, \tilde{M} . These lifts will be denoted $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{G}}$, respectively. We now give two descriptions of \bar{M} .

Let $[\rho] \in \mathcal{T}^2(\Sigma)$ be a (conjugacy class of) representation. Choose an identification of $\tilde{\Sigma}$ with \mathbb{H}^2 that is equivariant with respect to ρ , such an identification is called a *uniformization*. The uniformization allows us to equivariantly identify \bar{M} with the unit tangent bundle $S\mathbb{H}^2$. With this in mind we can think of a leaf of $\bar{\mathcal{G}}$ as an oriented geodesic in \mathbb{H}^2 and a leaf of $\bar{\mathcal{F}}$ as the union of all oriented geodesics with a common positive endpoint on $\partial\mathbb{H}^2$. Since




FIGURE 1. The identification between \overline{M} and $\partial\Gamma^{3+}$. Picture from [3]

$\mathrm{PSL}_2(\mathbb{R})$ acts simply transitively on $S\mathbb{H}^2$ we can further identify \overline{M} with $\mathrm{PSL}_2(\mathbb{R})$. Under this identification we see that $M = \rho(\Gamma)\backslash\mathrm{PSL}_2(\mathbb{R})$. Additionally, if we let

$$A = \left\{ \begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix} \right\}, \text{ and } P = \left\{ \begin{pmatrix} a & b \\ 0 & 1/a \end{pmatrix} \right\},$$

be *Cartan* and *parabolic* subgroups of $\mathrm{PSL}_2(\mathbb{R})$, then the leaves of the geodesic and stable foliations can be identified with right orbits of A and P , respectively.

The uniformization also allows us to identify the boundary of Γ , $\partial\Gamma$, with $\partial\mathbb{H}^2 \cong \mathbb{RP}^1$. Furthermore, this identification gives an action of Γ on $\partial\Gamma$ that is ρ -equivariant. This boundary gives us another way to describe \overline{M} , $\overline{\mathcal{F}}$, and $\overline{\mathcal{G}}$. $\partial\Gamma$ inherits an orientation coming from the orientation on \mathbb{H}^2 and we can identify \overline{M} with the set $\partial\Gamma^{3+}$ of pairwise distinct, positively oriented triples of points in $\partial\Gamma$. To see this observe that a point of \overline{M} can be thought of as an oriented geodesic, L , in \mathbb{H}^2 and a point, x , on L . Such a point can be identified with (t_+, t_0, t_-) , where t_+ and t_- are the positive and negative endpoints of L , respectively, and t_0 is the endpoint of the geodesic that intersects L perpendicularly and passes through x and makes the above triple positively oriented (see Figure 1). This identification is also equivariant with respect to the action of Γ and allows us to identify $\overline{\mathcal{F}}$ with $\partial\Gamma$ and $\overline{\mathcal{G}}$ with $\partial\Gamma^{(2)} := \partial\Gamma^2 \backslash \Delta$.

3. CONVEX FOLIATED PROJECTIVE STRUCTURES

In this section we describe the geometric structures whose moduli space $\mathcal{T}^4(\Sigma)$ describes. We begin with a description of projective structures. Roughly speaking, a projective structure on a manifold is a way to locally identify the manifold with \mathbb{RP}^n in such a way that the transition functions are locally elements of $\mathrm{PGL}_{n+1}(\mathbb{R})$. As such projective structures can be described in terms of atlases of charts. However, we will take a more global (but equivalent) point of view in our definition. Let N be an n -manifold. A *projective structure* consists of a pair (dev, hol) , where $hol : \pi_1(N) \rightarrow \mathrm{PGL}_{n+1}(\mathbb{R})$ is a representation and dev is a hol -equivariant local homeomorphism from \tilde{N} to \mathbb{RP}^n . Furthermore, we say that (dev_1, hol_1) and (dev_2, hol_2) are *equivalent* projective structures on N if there exists a homeomorphism $h : N \rightarrow N$ that is isotopic to the identity and an element $g \in \mathrm{PGL}_{n+1}(\mathbb{R})$ such that

- $dev_1 \circ \tilde{h} = g \circ dev_2$, where \tilde{h} is a lift of h to \tilde{N} , and
- $hol_2 = g^{-1}hol_1g$.

Let $\mathcal{P}(N)$ be the set of equivalence classes of projective structures on N . The above discussion shows that we have a map

$$hol : \mathcal{P}(N) \rightarrow \mathfrak{X}(\pi_1(N), \mathrm{PGL}_{n+1}(\mathbb{R})).$$

The map dev is called the *developing map* of the structure and the representation hol is called the *holonomy* of the structure.

As we mentioned before we are interested in projective structures that play well with the foliations coming from the geodesic flow. With this in mind, we say that a projective structure, (dev, hol) , on M is *foliated* if the following conditions are satisfied.

- For each leaf $\tilde{g} \in \tilde{\mathcal{G}}$, $dev(\tilde{g})$ is contained in a projective line, and
- For each leaf $\tilde{f} \in \tilde{\mathcal{F}}$, $dev(\tilde{f})$ is contained in a projective plane.

Two foliated projective structures are *equivalent* if they are equivalent as projective structures and the map $h : M \rightarrow M$ preserves the foliations \mathcal{F} and \mathcal{G} . We denote the set of equivalence classes of foliated projective structures by $\mathcal{P}_f(M)$. We now further refine this notion in order to arrive at the correct geometric structures. Let $C \subset \mathbb{RP}^n$, then C is *convex* if its intersection with every projective line is connected. If C is a convex subset of \mathbb{RP}^n then C is *properly convex* if its closure does not contain a affine line.

Exercise 3. *Show that a subset of \mathbb{RP}^n is properly convex if and only if its closure is contained in an affine patch.*

We can now define the appropriate projective structures. We say that a foliated projective structure on M is *convex* if the image of each leaf of $\tilde{\mathcal{F}}$ under the developing map is a convex set of a projective plane. Additionally, we define a *properly convex* foliated projective structure on M to be a foliated projective structure for which the image of each leaf of $\tilde{\mathcal{F}}$ is mapped to a properly convex subset of a projective plane by the developing map. Let $\mathcal{P}_{pcf}(M)$ subset of $\mathcal{P}_f(M)$ consisting of equivalence classes of properly convex foliated projective structures.

We can now rephrase the correspondence between $\mathcal{T}^4(\Sigma)$ and projective structures in more precise terms. Let $p : \bar{\Gamma} \rightarrow \Gamma$ be the projection implicit in (2.1). The map p gives an embedding of $\mathcal{T}^4(\Sigma) \subset \mathfrak{X}(\Gamma, \mathrm{PSL}_4(\mathbb{R})) \subset \mathfrak{X}(\bar{\Gamma}, \mathrm{PSL}_4(\mathbb{R}))$, and the correspondence can be succinctly stated as

Proposition 3.1. *The map hol is a homeomorphism between $\mathcal{P}_{pcf}(M)$ and $\mathcal{T}^4(\Sigma)$.*

Remark 3.2. *Since Γ has trivial center (2.1) implies that the center of $\bar{\Gamma}$ is cyclic, and we denote its generator by τ . The above correspondence implies that the holonomy of a properly convex foliated projective structure on M factors through p and thus every such holonomy kills τ .*

4. EXAMPLES AND IDEAS

In this section we will discuss certain examples of properly convex foliated projective structures on M and discuss some of the ideas required to prove Proposition 3.1. Let

$[\rho] \in \mathcal{T}^2(\Sigma)$, then we can define an element of $\mathcal{P}_{pcf}(M)$ as follows. Let $[Q] \in \mathbb{RP}^3$, where $Q = x(x^2 + y^2)$ (here we are using the fact that $\mathbb{R}^4 \cong \mathrm{Sym}^3(\mathbb{R}^2)$). Using the fact that $\overline{M} \cong \mathrm{PSL}_2(\mathbb{R})$ we can define a projective structure by letting dev^1 be the map $g \mapsto \rho_4(\rho(g)) \cdot [Q]$, where $g \in \mathrm{PSL}_2(\mathbb{R})$ and letting $hol = \rho_4 \circ \rho$. The vector $Q = (1, 0, 1, 0)$ in the standard basis for $\mathrm{Sym}^3(\mathbb{R}^2)$ and so

$$\begin{pmatrix} e^{t/2} & 0 \\ 0 & e^{-t/2} \end{pmatrix} \mapsto \begin{pmatrix} e^{3t/2} & 0 & 0 & 0 \\ 0 & e^{t/2} & 0 & 0 \\ 0 & 0 & e^{-t/2} & 0 \\ 0 & 0 & 0 & e^{-3t/2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} e^{3t/2} \\ 0 \\ e^{-t/2} \\ 0 \end{pmatrix} = \begin{pmatrix} e^{2t} \\ 0 \\ 1 \\ 0 \end{pmatrix},$$

and

$$\begin{pmatrix} a & b \\ 0 & 1/a \end{pmatrix} \mapsto \begin{pmatrix} a^3 & a^2b & ab^2 & b^3 \\ 0 & a & 2b & 3b^2/a \\ 0 & 0 & 1/a & 3b/a^2 \\ 0 & 0 & 0 & 1/a^3 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} a^3 + ab^2 \\ 2b \\ 1/a \\ 0 \end{pmatrix} = \begin{pmatrix} a^2(a^2 + b^2) \\ 2ab \\ 1 \\ 0 \end{pmatrix}.$$

Under the coordinate change $v = a^2(a^2 + b^2)$, $u = 2ab$, we see that the above image can be identified with $v > u^2/4$, and is thus properly convex (See Exercise 3). In fact, this set can be identified with \mathbb{H}^2 (see Exercise 2). Thus we see that this projective structure is properly convex foliated.

Despite knowing that these structures are properly convex foliated, we do not have a very good idea of what they look like globally. In order to get a more global picture we will try to understand the developing map in terms of the description of \overline{M} as $\partial\Gamma^{3+}$. Let V be a vector space and let $Flag(V)$ denote the flag variety of V . If we think of \mathbb{R}^2 as $\mathrm{Sym}^1(\mathbb{R}^2)$, then the *Veronese embedding* gives the following equivariant curve $\xi : \partial\Gamma \cong \mathbb{RP}^1 \rightarrow Flag(\mathbb{R}^4)$. Given by $\xi = (\xi^1, \xi^2, \xi^3)$, where

- $\xi^1([S])$ is the line of polynomials divisible by S^3 ,
- $\xi^2([S])$ is the plane of polynomials divisible by S^2 , and
- $\xi^3([S])$ is the hyperplane of polynomials divisible by S .

If we let Ω_ξ be the set of polynomials in \mathbb{RP}^3 with a single real root (i.e. they factor over \mathbb{R} into a linear and a quadratic term), then Ω_ξ is the image of dev .

Exercise 4. *Prove that Ω_ξ is the image of dev , namely that Ω_ξ is the $\mathrm{PSL}_2(\mathbb{R})$ orbit of $[Q]$ under the action $g \cdot [R] = \rho_4(\rho(g)) \cdot [R]$.*

The map ξ allows us to define a family, ξ_t^1 of equivariant maps from $\partial\Gamma \rightarrow \mathbb{RP}^3$, given by

$$\xi_t^1(t') = \begin{cases} \xi^3(t) \cap \xi^2(t') & \text{if } t \neq t' \\ \xi^1(t) & \text{if } t = t' \end{cases}$$

For each t , the image of ξ_t^1 in $\xi^3(t)$ bounds the copy of \mathbb{H}_t^2 given by $dev(t)$ (here we are thinking of t as a leaf of $\overline{\mathcal{F}}$). The geodesic leaf $g = (t_+, t_-)$ is taken to the intersection of \mathbb{H}_t^2 and the projective line, $\overline{\xi^1(t_+) \xi_{t_+}^1(t_-)}$, connecting $\xi^1(t_+)$ and $\xi_{t_+}^1(t_-)$. The tangent lines to \mathbb{H}_t^2 at $\xi^1(t_+)$ and $\xi_{t_+}^1(t_-)$ are $\xi^2(t_+)$ and $\xi^3(t_+) \cap \xi^3(t_-)$, respectively. Furthermore, these

¹Technically, dev is a lift of this map to \tilde{M} .




FIGURE 2. The image of the developing map. Picture from [3].

lines intersect in the point $\xi^3(t_-) \cap \xi^2(t_+) = \xi^1_{t_+}(t_-)$. Finally, given t_0 different from t_+ and t_- there is a unique point of intersection between $\overline{\xi^1(t_+)\xi^1_{t_+}(t_-)}$ and $\overline{\xi^1_{t_+}(t_+)\xi^1_{t_+}(t_0)}$. Thus we see that in terms of $\partial\Gamma^{3+}$ that dev is defined by

$$(4.1) \quad (t_+, t_0, t_-) \mapsto \overline{\xi^1(t_+)\xi^1_{t_+}(t_-)} \cap \overline{\xi^1_{t_+}(t_+)\xi^1_{t_+}(t_0)}.$$

The discussion of the previous paragraph is illustrated in Figure 2

5. CONVEX REPRESENTATIONS

The proof of Proposition 3.1 relies on the fact that representations in $\mathcal{T}^4(\Sigma)$ can be characterized by a certain convexity property which we now discuss. Work of Labourie and Guichard [5, 7] has shown that $\rho \in \mathcal{T}^4(\Sigma)$ if and only if ρ is a convex representation. A representation is *convex* if we can find a ρ -equivariant curve $\xi^1 : \partial\Gamma \rightarrow \mathbb{RP}^3$ such that for t_1, \dots, t_4 that are pairwise distinct,

$$\xi^1(t_1) \oplus \dots \oplus \xi^1(t_4) = \mathbb{R}^4.$$

A simple exercise shows that a convex curve in \mathbb{RP}^2 bounds a properly convex set. Additionally, a curve $\xi = (\xi^1, \dots, \xi^{n-1}) : \mathbb{RP}^1 \rightarrow Flag(\mathbb{R}^n)$ is called *Frenet* (this is sometimes known as hyperconvex) if

- (1) For every (n_1, \dots, n_k) such that $\sum_{i=1}^k n_i = n$ and every $t_1, \dots, t_k \in \mathbb{RP}^1$ of pairwise distinct elements the following sum is direct

$$\sum_{i=1}^k \xi^{n_i}(x_i) = \mathbb{R}^n.$$

(2) For every (m_1, \dots, m_k) with $\sum_{i=1}^k m_i = m \leq n$ and every $x \in \mathbb{RP}^1$

$$\lim_{(x_i) \rightarrow x} \sum_{i=1}^k \xi^{m_i}(x_i) = \xi^m(x),$$

where the limit is taken over k -tuples of pairwise distinct points.

It is easy to see that if ξ is Frenet then ξ^1 is a convex curve and that ξ^1 determines ξ via the limit condition. By work of Labourie [7] has shown that If a ρ is convex then it is possible to find a unique ρ -equivariant Frenet curve. Given a non-degenerate bilinear form on \mathbb{R}^n and a curve $f = (f^1, \dots, f^{n-1}) : \mathbb{RP}^1 \rightarrow \mathit{Flag}(\mathbb{R}^n)$, it is possible to define a dual curve $f^\perp = (f^{n-1,\perp}, \dots, f^{1,\perp}) : \mathbb{RP}^1 \rightarrow \mathit{Flag}(\mathbb{R}^{n*})$. Work of Guichard [4] shows that a curve ξ is Frenet if and only if ξ^\perp is Frenet. This duality will be crucial in the proof of Proposition 3.1.

5.1. convex implies properly convex foliated. We begin by showing that an element $[\rho] \in \mathcal{T}^4(\Sigma)$ gives rise to a properly convex foliated projective structure on M . By the previous paragraph we see that ρ is a convex representation and thus we can find a ρ -equivariant flag curve ξ . We begin by using this curve to define a family of ρ -equivariant lower dimensional flag curves. Define $\xi_t : \mathbb{RP}^1 \rightarrow \mathit{Flag}(\xi^3(t))$ by

$$(5.1) \quad \xi_t(t') = \begin{cases} (\xi^3(t) \cap \xi^2(t'), \xi^3(t) \cap \xi^3(t')) & \text{if } t \neq t' \\ (\xi^1(t), \xi^2(t)) & \text{if } t = t' \end{cases}$$

For each $t \in \mathbb{RP}^1$ the curve ξ_t is also Frenet. This is proven by showing that ξ_t^\perp is Frenet and using the basic fact that if W and V are linear subspace then $(V + W)^\perp = V^\perp \cap W^\perp$. For example, to show that ξ_t^\perp is Frenet one of the things we need to show that $(\xi_t^{2,\perp}(t_1) + \xi_t^{1,\perp}(t_2)) = \mathbb{R}^3$ for all distinct pairs t_1, t_2 . To show this we observe that

$$(\xi_t^{2,\perp}(t_1) + \xi_t^{1,\perp}(t_2))^\perp = \xi_t^2(t_1) \cap \xi_t^1(t_2) = \xi^3(t) \cap \xi^3(t_1) \cap \xi^2(t_2) = (\xi^{3,\perp}(t) + \xi^{3,\perp}(t_1) + \xi^{2,\perp}(t_2))^\perp = \{0\},$$

with the last equality coming from the fact that ξ^\perp is Frenet. The other conditions needed to be verified to show that ξ^\perp is Frenet can all be shown in a similar way. We now define a developing map using the formula in (4.1). For each t the image of ξ_t^\perp bounds is convex and thus bounds a properly convex subset C_t of $\xi^3(t)$. The Frenet properties of ξ this new developing map has all the same nice properties as the map given to us in the previous example by the Veronese embedding.

5.2. properly convex foliated implies convex. The more difficult direction is to show that given a properly convex foliated projective structure on M that the holonomy representation is a convex representation. Details can be found in [3] and we simply outline the key ideas. Suppose that we have such a structure with holonomy ρ . Since we know that the image of a leaf of $\tilde{\mathcal{F}} \cong \tilde{\Gamma}$ under the developing map is contained in a projective plane we get a map² $\xi^3 : \partial\tilde{\Gamma} \rightarrow \mathbb{RP}^{3*}$ taking $t \in \partial\tilde{\Gamma}$ to the projective plane containing $dev(t)$. The first thing that we have to do is to show that the map ξ^3 descends to a map defined on $\partial\Gamma$ (this

²Here we are implicitly identifying the space of projective planes in \mathbb{RP}^3 with \mathbb{RP}^{3*} using a non-degenerate bilinear form.

part is highly non-trivial and comprises a good chunk of [3]). We must then show that this map is convex. Namely that for t_1, \dots, t_4 pairwise distinct points that

$$\xi^3(t_1) + \dots + \xi^3(t_4) = \mathbb{R}^{4*},$$

or equivalently that

$$(5.2) \quad \xi^3(t_1) \cap \dots \cap \xi^3(t_4) = \emptyset.$$

The fact that these planes do not have a common intersection can be viewed geometrically. Fix t_1 , then the fact that the intersection from (5.2) is empty is equivalent to the three lines $\xi^3(t_1) \cap \xi^3(t_i)$, $2 \leq i \leq 4$, not intersecting. Let C_{t_1} be the properly convex set that is the image of the developing map restricted to plane $\xi^3(t_1)$. Then it can be shown (with a good deal of work) that the domain C_t is *strictly convex* (contains no line segments in its boundary) and that the lines $\xi^3(t_1) \cap \xi^3(t_i)$ for $2 \leq i \leq 4$ are tangent lines to C_{t_1} at distinct points, and thus do not intersect. Try drawing tangents to the domain in Figure 2 to convince yourself that these lines must be disjoint.

REFERENCES

- [1] Suhyoung Choi and William M. Goldman, *Convex real projective structures on closed surfaces are closed*, Proc. Amer. Math. Soc. **118** (1993), no. 2, 657–661. MR1145415 (93g:57017)
- [2] William M. Goldman, *Convex real projective structures on compact surfaces*, J. Differential Geom. **31** (1990), no. 3, 791–845. MR1053346 (91b:57001)
- [3] Olivier Guichard, *Une dualité pour les courbes hyperconvexes*, Geom. Dedicata **112** (2005), 141–164. MR2163895 (2007a:53103)
- [4] ———, *Composantes de Hitchin et représentations hyperconvexes de groupes de surface*, J. Differential Geom. **80** (2008), no. 3, 391–431. MR2472478 (2009h:57031)
- [5] Olivier Guichard and Anna Wienhard, *Convex foliated projective structures and the Hitchin component for $\mathrm{PSL}_4(\mathbf{R})$* , Duke Math. J. **144** (2008), no. 3, 381–445. MR2444302 (2009k:53223)
- [6] N. J. Hitchin, *Lie groups and Teichmüller space*, Topology **31** (1992), no. 3, 449–473. MR1174252 (93e:32023)
- [7] François Labourie, *Anosov flows, surface groups and curves in projective space*, Invent. Math. **165** (2006), no. 1, 51–114. MR2221137 (2007c:20101)